



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\frac{du}{dC_1} = \frac{C}{1+C^2C_1^2}, \quad \frac{du}{dC_2} = -\frac{C}{1+C^2C_2^2},$$

$$\frac{d^2u}{dC_1^2} = -\frac{2C^3C_1}{(1+C^2C_1^2)^2}, \quad \frac{d^2u}{dC_2^2} = \frac{2C^3C_2}{(1+C^2C_2^2)^2}, \quad \frac{d^2u}{dC_1dC_2} = 0.$$

$$\text{Now } \frac{d^2u}{dC_1^2} \frac{d^2u}{dC_2^2} < \frac{d^2u}{dC_1dC_2}.$$

Hence there is no maximum unless  $C_1 = \infty$ ,  $C_2 = -\infty$ .

**II. Solution by V. M. SPUNAR, Chicago, Ill., and the PROPOSER.**

By the principle of the instrument, we have

$$\frac{\tan\phi_1}{\tan\phi_2} = \frac{c_1}{c_2} \quad \therefore \frac{\tan\phi_1 - \tan\phi_2}{\tan\phi_1 + \tan\phi_2} = \frac{c_1 - c_2}{c_1 + c_2} \dots (1).$$

But by trigonometry,

$$\tan\phi_1 - \tan\phi_2 = \frac{\sin(\phi_1 - \phi_2)}{\cos\phi_1 \cos\phi_2}, \quad \tan\phi_1 + \tan\phi_2 = \frac{\sin(\phi_1 + \phi_2)}{\cos\phi_1 \cos\phi_2}.$$

Hence (1) becomes

$$\frac{\sin(\phi_1 - \phi_2)}{\sin(\phi_1 + \phi_2)} = \frac{c_1 - c_2}{c_1 + c_2}, \text{ or } \sin(\phi_1 - \phi_2) = \left( \frac{c_1 - c_2}{c_1 + c_2} \right) \sin(\phi_1 + \phi_2).$$

Hence,  $(\phi_1 - \phi_2)$  is a maximum when  $\sin(\phi_1 + \phi_2)$  is maximum; that is, when  $\sin(\phi_1 + \phi_2) = 1$ ; that is,  $\phi_1 + \phi_2 = \frac{1}{2}\pi$ .

**297. Proposed by PROF. L. C. WALKER, Socorro, New Mexico.**

A square hole  $2s$  on a side is cut through an ellipsoid, axes  $2a$ ,  $2b$ ,  $2c$ , the axis of the hole coinciding with the axis  $2c$  of the ellipsoid. Find (1) the volume, and (2) the surface removed.

**Solution by J. SCHEFFER, A. M., Hagerstown, Md.**

$$\text{The volume } V = 8c \int_0^s dx \int_0^s dy \sqrt{1 - \left(\frac{x}{a}\right)^2 - \left(\frac{y}{b}\right)^2}$$

$$\begin{aligned}
&= \frac{4c}{b} \left[ \frac{bs}{a} \int_0^s \sqrt{\frac{a^2(b^2-s^2)}{b^2}} - x^2 + \frac{b^2}{a^2} \int_0^s (a^2-x^2) \sin^{-1} \frac{as}{b \sqrt{(a^2-x^2)}} dx \right. \\
&= \frac{4bcs}{3a^2(3a^2-s^2)} \sin^{-1} \frac{as}{b \sqrt{(a^2-s^2)}} + \frac{4acs(3b^2-s^2)}{3b^2} \sin^{-1} \frac{bs}{a \sqrt{(b^2-s^2)}} \\
&\quad \left. + \frac{8}{3} abcs \sin^{-1} \frac{s^2}{(a^2-s^2)^{\frac{1}{2}}(b^2-s^2)^{\frac{1}{2}}} + \frac{10cs^2}{3ab} \sqrt{[a^2b^2-(a^2+b^2)s^2]} \right].
\end{aligned}$$

The surface  $S = \frac{4}{ab} \int_0^s dx \int_0^s dy \sqrt{\frac{a^4b^4 - b^4(a^2-c^2)x^2 - a^4(b^2-c^2)y^2}{a^2b^2 - b^2x^2 - a^2y^2}}.$

The integration of this leads to elliptical functions.

NOTE. If any one will complete the solution of the second part of this problem we shall be pleased to publish it. ED. F.

## NUMBER THEORY AND DIOPHANTINE ANALYSIS.

NOTE. Dr. Whalen, of the University of Illinois, has consented to edit this department in the future. Will contributors please send their problems and solutions to him?

174. Proposed by B. KRAMER, Student, University of Pittsburg, Pittsburg, Pa.

Find a general solution of  $x(x+a)=y^2$ ,  $a$ ,  $x$ , and  $y$  being integers. Given  $a$ , required to find  $x$  to satisfy conditions.

I. Solution by PROFESSOR F. L. GRIFFIN, Ph. D., Williams College.

Let  $d$  be any integer contained in  $a$  an odd number of times. Then the general solution is:  $x=(a-d)^2/4d$ ,  $y=(a^2-d^2)/4d$ .

For let  $y=x+n$ , whence  $x^2+ax=x^2+2nx+n^2$ , or  $x=n^2/(a-2n)$ .

Now  $x$  is integral if, and only if,  $a-2n$  is a factor of  $n$ ; that is, if an integer  $k$  exists such that  $n=kd$  [denoting  $a-2n$  by  $d$ ]. But then  $a=d+2n=d(1+2k)$ ; so that integral values of  $x$  exist if, and only if,  $d$  is contained in  $a$  an odd number of times.

Using any such number  $d$  we have,  $k=(a-d)/2d$ ,  $n=\frac{1}{2}(a-d)$ ,  $x=(a-d)^2/4d$ ,  $y=x+\frac{1}{2}(a-d)=(a^2-d^2)/4d$ , as stated.

Remarks. (I) Negative values of  $d$  are admissible, as the factors  $x$  and  $x+a$  merely have their numerical values interchanged.

(II) If  $a$  be prime,  $d=\pm 1$  gives the only solution except the trivial case  $d=a$ ,  $x=y=0$ .

(III) Numerical examples follow: